

Corrections

In [1], the Figure 2 legend should be amended from:

Figure 2 Ultrasound screen captures of block administration, with needle advanced cranio-caudally. 1 ribs, 2 serratus anterior muscle, 3 intercostal muscle, 4 local anaesthetic.

to:

Figure 2 Ultrasound screen captures of block administration, with needle advanced cranio-caudally. 1 paraspinal muscles, 2 local anaesthetic, 3 intercostal muscles, 4 ribs.

Reference

1. Roué C, Wallaert M, Kacha M, Havet E. Intercostal/paraspinal nerve block for thoracic surgery. *Anaesthesia* 2015; **71**: 112–3.

doi:10.1111/anae.13383

In the Guidelines [1] by Harrop-Griffiths et al., the notes to accompany Table 2 stated that there was one death in a patient who underwent lumbar plexus block. This was incorrect, and should read:

“The one death in this series was that of a patient on clopidogrel who underwent a lumbar sympathetic block. . .”

Reference

1. Harrop-Griffiths W, Cook T, Gill H, Hill D, Ingram M, Makris M, Malhotra S, Nicholls B, Popat M, Swales H, Wood P (Working Party). The Association of Anaesthetists of Great Britain & Ireland & The Obstetric Anaesthetists' Association Regional Anaesthesia UK. Regional anaesthesia and patients with abnormalities of coagulation. *Anaesthesia* 2013; **68**: 966–72.

doi:10.1111/anae.13380